



***In silico* reconstruction of protein-protein interaction networks of bacteria used in synthetic biology**

C.T. Chasapis^{1#}, V. Savvopoulou² and M.I. Klapa^{2*}

¹ Metabolic Engineering & Systems Biology Laboratory, FORTH/ICE-HT, Patras, Greece

² Graduate Program "Chemical Biology", Department of Chemistry, University of Patras, Patras, Greece

Presenting author: C.T. Chasapis, cchasapis@upatras.gr

* Corresponding author: M.I. Klapa, mklapa@iceht.forth.gr

ABSTRACT

The genetic and physiological characterization of bacteria that are metabolic engineering and synthetic biology resources is essential for the identification of appropriate targets for genetic modification towards the development of more efficient strains and the optimization of their fermentation processes. The number of fully sequenced bacteria has dramatically increased in recent years, providing a valuable basis for the design of high-throughput biomolecular (omic) analyses that can help elucidate the regulatory machinery connecting genotype to phenotype. The interpretation of physiological and omic measurement datasets could be greatly assisted by the accurate reconstruction and integrated analysis of the metabolic and protein-protein interaction (PPI) networks of the investigated bacteria. While a lot of effort has been placed in the genome-scale metabolic network reconstruction in bacteria, being a focal area of research in metabolic engineering for many years, this is not the case with the bacterial PPI networks. To-date, experimentally reconstructed PPI networks based on high-throughput experiments exist for very few bacterial systems. Thus, for most of the bacteria, there is a need to *in silico* reconstruct their potential PPI network based on any available experimental data and bioinformatic resources using predictive models.

In light of this limited experimental PPI data availability for microbial systems, we designed and applied a workflow for the *in silico* reconstruction of the PPI network of two bacteria, *Streptomyces lividans* [1] and *Moorella thermoacetica* [2], which have been used in synthetic biology applications in the recent years. *S. lividans* is a preferred host for the overproduction of secretory proteins and secondary metabolites of industrial and pharmaceutical interest, however, it has a very small number of experimentally determined PPIs reported in source databases. Through the applied workflow, we were able to identify more than a thousand experimentally supported PPIs based on comparative genomic analysis with evolutionarily adjacent bacterial systems. *M. thermoacetica* is a thermophilic Gram-positive acetogen that has been used for the non-photosynthetic CO₂ fixation for the production of useful chemicals. While there is a limited number of experimentally determined PPIs for this bacterium too, a larger than for *S. lividans* number of multi-omic datasets are available, supporting a more reliable application of bioinformatics predictive algorithms. Our analysis lead to the reconstruction of a network of 12,910 PPIs between 2,112 proteins, which can be used to enrich our understanding of the molecular physiology of this microorganism, in conjunction with the available metabolic network. The reconstructed PPI networks of both microbes can form the basis for validation studies of key PPIs and be used in the design of synthetic biology experiments.

"This work was supported by the EC 7th Framework Programme STREPSYNTH project (n° 613877), the NSRF 2014-2020 ELIXIR-GR (MIS 5002780) and INSPIRED (5002550) projects, co-financed by Greece and the European Union (European Regional Development Fund) and the BIOMEK (T1EAK-00279) project, co-financed by Greece and the European Union through the Operational Program Competitiveness, Entrepreneurship and Innovation, under the call RESEARCH – CREATE – INNOVATE."

REFERENCES

- [1] Chasapis CT, Rückert C, Busche T, Kalinowski J, Economou A, Klapa MI. (2019) The *in silico* reconstruction of the *Streptomyces lividans* protein-protein interaction network (in prep).
- [2] Savvopoulou V. (2019) *In silico* reconstruction of the protein-protein interaction network of *Moorella thermoacetica*, used for the non-photosynthetic CO₂ bioconversion into useful chemicals. Master's Thesis, U. Patras (Supervisors: A. Vlamis & M.I. Klapa).